

Question 1

$$(a) \quad \frac{1+i}{2-i} = \frac{1+i}{2-i} \frac{2+i}{2+i} = \frac{1+3i}{5}. \quad (\text{A1, Sect. 1, Para. 6})$$


$$(b) \quad -i = \cos\left(-\frac{\pi}{2}\right) + i \sin\left(-\frac{\pi}{2}\right)$$

The principal cube root is (A1, Sect. 3, Para. 3)

$$\cos\left(\frac{1}{3}\left(-\frac{\pi}{2}\right)\right) + i \sin\left(\frac{1}{3}\left(-\frac{\pi}{2}\right)\right) = \cos\left(\frac{\pi}{6}\right) - i \sin\left(\frac{\pi}{6}\right) = \frac{\sqrt{3}}{2} - i \frac{1}{2}.$$

$$(c) \quad \text{Log}\left(\frac{1+i}{\sqrt{2}}\right) = \log_e\left(\left|\frac{1+i}{\sqrt{2}}\right|\right) + i \text{Arg}\left(\frac{1+i}{\sqrt{2}}\right) = \log_e 1 + \frac{\pi}{4}i = \frac{\pi}{4}i \quad (\text{A2, Sect. 5, Para. 1})$$

$$(d) \quad \begin{aligned} \left(\frac{1+i}{\sqrt{2}}\right)^{2-4i} &= \exp\left((2-4i) \text{Log}\left(\frac{1+i}{\sqrt{2}}\right)\right) && (\text{A2, Sect. 5, Para. 3}) \\ &= \exp\left((2-4i)\frac{\pi}{4}i\right) && \text{using the result from part (c)} \\ &= \exp\left(i\frac{\pi}{2}\right) \exp(\pi) = i \exp(\pi). \end{aligned}$$

Question 2**(a)****(b)(i)** A and A - B are regions. (A3, Sect. 4, Para. 6)

[[B is not a region as it is not open.]]

(b)(ii) A - B is a simply-connected region. (B2, Sect 1, Para. 3)[[In A the inside of $|z - i| = 1.5$ contains points not in A]]**(b)(iii)** B is closed. (A3, Sect. 5, Para. 1)**(b)(iv)** B is compact. (A3, Sect. 5, Para.5)

Question 3

(a)

(a)(i) The standard parametrization for the line segment Γ is (A2, Sect. 2, Para. 3)

$$\gamma(t) = (1-t)i + t(1-i) = t + i(1-2t) \quad (t \in [0, 1])$$

(a)(ii) Since γ is a smooth path then (B1, Sect. 2, Para. 1)

$$\begin{aligned} \int_{\Gamma} \operatorname{Re} z \, dz &= \int_0^1 \operatorname{Re}(\gamma(t)) \gamma'(t) \, dt \\ &= \int_0^1 t(1-2i) \, dt = (1-2i) \left[\frac{t^2}{2} \right]_0^1 = \frac{1-2i}{2} \end{aligned}$$

(b)

The length of Γ is $L = |(1-i) - i| = |1-2i| = \sqrt{5}$.Using the Triangle Inequality (A1, Sect. 5, Para. 3b) then, for $z \in \Gamma$, we have

$$\begin{aligned} |\cos z| &= \frac{1}{2} |e^{iz} + e^{-iz}| \leq \frac{1}{2} (|e^{iz}| + |e^{-iz}|) = \frac{1}{2} \{ e^{\operatorname{Re}(iz)} + e^{\operatorname{Re}(-iz)} \} \quad (\text{A2, Sect. 4, Para. 2b}) \\ &= \frac{1}{2} \{ e^{-y} + e^y \} \leq \frac{1}{2} \{ e^1 + e^1 \} = e \end{aligned}$$

Using the Backwards form of the Triangle Inequality (A1, Sect. 5, Para. 3c) then, for $z \in \Gamma$, we have

$$|6+z^2| \geq \left| |6| - |z^2| \right| \geq |6 - |1+i|^2| = |6-2| = 4.$$

Therefore $M = \left| \frac{\cos z}{6+z^2} \right| \leq \frac{e}{4}$ for $z \in \Gamma$. $f(z) = \frac{\cos z}{6+z^2}$ is continuous on $\mathbb{C} - \{-\sqrt{6}i, \sqrt{6}i\}$ and hence on the line Γ .

Therefore by the Estimation Theorem (B1, Sect. 4, Para. 3)

$$\left| \int_{\Gamma} f(z) \, dz \right| \leq ML = \frac{e}{4} \sqrt{5}.$$

Question 4

(a)

Let \mathcal{R} be the simply-connected region $\{z: |z| < 2\}$. C is a closed contour in \mathcal{R} , and $\frac{\cos z}{(z-3)^3}$ is analytic on \mathcal{R} .

By Cauchy's Theorem (B2, Sect. 1, Para. 4)

$$\int_C \frac{\cos z}{(z-3)^3} dz = 0.$$

(b)

Let \mathcal{R} be the simply connected region $\{z: |z| < 2\}$. \mathcal{R} is a simply-connected region and C is a simple-closed contour in \mathcal{R} . As $f(z) = \frac{\cos z}{(z-3)^2}$ is analytic on \mathcal{R} then by Cauchy's Integral Formula (B2, Sect. 2, Para. 1)

$$\int_C \frac{\cos z}{z(z-3)^2} dz = \int_C \frac{f(z)}{z-0} dz = 2\pi i f(0) = 2\pi i \frac{\cos 0}{(-3)^2} = \frac{2}{9} \pi i.$$

(c)

Let \mathcal{R} be the simply connected region \mathbb{C} . \mathcal{R} is a simply-connected region and C is a simple-closed contour in \mathcal{R} . As $f(z) = \cos z$ is analytic on \mathcal{R} then by Cauchy's n'th Derivative Formula (B2, Sect. 3, Para. 1) with $n = 2$ and $\alpha = 0$ we have

$$\int_C \frac{\cos z}{z^3} dz = \frac{2\pi i}{2!} f^{(2)}(0) = \pi i (-\cos 0) = -\pi i.$$

Question 5

(a)

$f(z) = \frac{1}{2(z + \frac{1}{2})(z + 2)}$. Therefore z has simple poles at $z = -1/2$ and $z = -2$.

$$\text{Res}(f, -\frac{1}{2}) = \lim_{z \rightarrow -\frac{1}{2}} (z + \frac{1}{2}) f(z) = \frac{1}{2(-\frac{1}{2} + 2)} = \frac{1}{3}. \quad [[\text{C1, Sect. 1, Para. 1}]]$$

$$\text{Res}(f, -2) = \lim_{z \rightarrow -2} (z + 2) f(z) = \frac{1}{2(-2 + \frac{1}{2})} = -\frac{1}{3}.$$

(b)

I shall use the strategy given in C1, Sect. 2, Para. 2.

$$\int_0^{2\pi} \frac{1}{5 + 4 \cos t} dt = \int_C \frac{1}{5 + 4 \frac{1}{2}(z + z^{-1})} \frac{1}{iz} dz, \quad \text{where } C \text{ is the unit circle.}$$

$$= -i \int_C \frac{1}{2z^2 + 5z + 2} dz = -i \int_C \frac{1}{(2z + 1)(z + 2)} dz.$$

The only residue inside the unit circle is the one at $z = -1/2$.

$$\text{Therefore } \int_0^{2\pi} \frac{1}{5 + 4 \cos t} dt = -i * 2\pi i \text{Res}(f, -\frac{1}{2}) = \frac{2}{3} \pi.$$

[[As it is a real integral we expect the answer will be real.]]

Question 6

(a)(i) Let $g_1(z) = z^5$.

For $z \in C_1$ then, using the Triangle Inequality (A1, Sect. 5, Para. 3),

$$|f(z) - g_1(z)| = |3z^2 + i| \leq |3z^2| + |i| = 12 + 1 < 32 = |g_1(z)|.$$

As f is a polynomial then it is analytic on the simply-connected region $\mathbf{R} = \mathbf{C}$. Since C_1 is a simple-closed contour in \mathbf{R} then by Rouché's theorem (C2, Sect. 2, Para. 4) f has the same number of zeros as g_1 inside the contour C_1 . Therefore f has 5 zeros inside C_1 .

(a)(ii) Let $g_2(z) = 3z^2$.

On the contour C_2 we have, using the Triangle Inequality,

$$|f(z) - g_2(z)| = |z^5 + i| \leq |z^5| + |i| = 1 + 1 < 3 = |g_2(z)|.$$

As C_2 is a simple-closed contour in \mathbf{R} then by Rouché's theorem f has the same number of zeros as g_2 inside the contour C_2 . Therefore f has 2 zeros inside C_2 .

(b)

From part(a) $f(z)$ has $5 - 2 = 3$ solutions in the set $\{z: 1 \leq z < 2\}$. Therefore we have to find if there are any solutions on C_2 .

From part (a), on C_2 we have $|z^5 + i| \leq 2$.

Therefore, using the Backwards form of the Triangle Inequality (A1, Sect. 5, Para. 3c)

$$|f(z)| \geq | |3z^3| - |z^5 + i| | \geq |3 - 2| = 1, \text{ on } C_2.$$

As $f(z)$ is non-zero on C_2 then there are exactly 3 solutions in the set $\{z: 1 < z < 2\}$.

Question 7

(a)

The conjugate velocity function $\bar{q}(z) = 2/z$.

As q is a steady continuous 2-dimensional velocity function on the region $\mathbb{C} - \{0\}$ and \bar{q} is analytic on $\mathbb{C} - \{0\}$ then q is a model fluid flow (D2, Sect. 1, Para. 14).

(b) On $\mathbb{C} - \{0\}$, $\Omega(z) = 2 \text{Log } z$ is a primitive of \bar{q} . Therefore Ω is a complex potential function for the flow (D2, Sect. 2, Para. 1).

The stream function $\Psi(x, y) = \text{Im} \Omega(z)$ (D2, Sect. 2, Para. 4)
 $= 2 \text{Im}(\log_e |z| + i \text{Arg } z) = 2 \text{Arg } z$.

A streamline through the point i satisfies the equation


$$2 \text{Arg } z = \Psi(0,1) = 2 \frac{\pi}{2}. \text{ So } \text{Arg } z = \frac{\pi}{2}. \quad (\text{D2, Sect. 2, Para. 4})$$

A streamline through the point $1 + i$ satisfies the equation

$$2 \text{Arg } z = \Psi(1,1) = 2 \frac{\pi}{4}. \text{ So } \text{Arg } z = \frac{\pi}{4}.$$

Since $q(i) = \frac{2}{-i} = 2i$ the flow at i is in the y direction.

As $q(1+i) = \frac{2}{1-i} = 1+i$ the flow at $1+i$ is in the North-East direction.



(c)

The flux of q across the unit circle $C = \{z : |z| = 1\}$ is (D2, Sect. 1, Para. 10)

$$\text{Im} \left(\int_C \bar{q}(z) dz \right) = \text{Im} \left(\int_C \frac{2}{z} dz \right) = \text{Im}(2 * 2\pi i) = 4\pi \quad \text{, using Cauchy's Integral Theorem}$$

(B2, Sect. 2, Para. 1).

Question 8

(a)

If α is a fixed point of f then $f(\alpha) = \alpha^2 - 4\alpha + 6 = \alpha$ (D3, Sect. 1, Para 3).

As $\alpha^2 - 5\alpha + 6 = (\alpha - 2)(\alpha - 3) = 0$ then there are fixed points at 2 and 3.

$$f'(z) = 2z - 4.$$

As $f'(2) = 0$ then 2 is a super-attracting fixed point (D3, Sect. 1, Para. 5).

As $f'(3) = 2$ then 3 is a repelling fixed point.

(b)(i) [[From the diagram in Handbook looks as if point not in Mandelbrot set.]]

$$P_c(0) = -1 + i.$$

$$P_c^2(0) = (-1 + i)^2 + (-1 + i) = -1 - i.$$

$$P_c^3(0) = (-1 - i)^2 + (-1 + i) = -1 + 3i.$$

As $|P_c^3(0)| > 2$ then c does not lie in the Mandelbrot set (D3, Sect. 4, Para. 5).

(b)(ii)

Since $|c+1| = \left|\frac{1}{8}i\right| < \frac{1}{4}$ then P_c has an attracting 2-cycle (D3, Sect. 4, Para. 9(b)).

Therefore c belongs to the Mandelbrot set (D3, Sect. 4, Para. 8).

Question 9

(a)

(a)(i)

$$f(z) = \bar{z} \operatorname{Im} z + |z|^2 = (x - iy)y + (x^2 + y^2) = u(x, y) + iv(x, y),$$

where $u(x, y) = x^2 + xy + y^2$, and $v(x, y) = -y^2$.

(a)(ii)

$$\frac{\partial u}{\partial x} = 2x + y, \quad \frac{\partial u}{\partial y} = x + 2y, \quad \frac{\partial v}{\partial x} = 0, \quad \frac{\partial v}{\partial y} = -2y$$

If f is differentiable then the Cauchy-Riemann equations hold (A4, Sect. 2, Para. 1). If they hold at (a, b) then

$$\frac{\partial u}{\partial x}(a, b) = 2a + b = -2b = \frac{\partial v}{\partial y}(a, b), \text{ and}$$

$$\frac{\partial v}{\partial x}(a, b) = 0 = -(a + 2b) = -\frac{\partial u}{\partial y}(a, b)$$

Therefore the Cauchy-Riemann equations only hold at $(0, 0)$.

As f is defined on the region \mathbf{C} , and the partial derivatives $\frac{\partial u}{\partial x}, \frac{\partial u}{\partial y}, \frac{\partial v}{\partial x}, \frac{\partial v}{\partial y}$

1. exist on \mathbf{C}
2. are continuous at $(0, 0)$.
3. satisfy the Cauchy-Riemann equations at $(0, 0)$

then, by the Cauchy-Riemann Converse Theorem (A4, Sect. 2, Para. 3), f is differentiable at 0.

As the Cauchy-Riemann equations only hold at $(0, 0)$ then f is not differentiable on any region surrounding 0. Therefore f is not analytic at 0. (A4, Sect. 1, Para. 3)

(a)(iii)



$$f'(0, 0) = \frac{\partial u}{\partial x}(0, 0) + i \frac{\partial v}{\partial x}(0, 0) = 0 \quad (\text{A4, Sect. 2, Para. 3}).$$

(b)(i)

The domain of g is \mathbf{C} (A4, Sect. 1, Para. 7) and its derivative $g'(z) = 3z^2$ also has domain \mathbf{C} (A4, Section 3, Para. 4). Therefore g is analytic on \mathbf{C} . Since $g'(z) \neq 0$ when $z \neq 0$ then g is conformal on $\mathbf{C} - \{0\}$ (A4, Sect. 4, Para. 6).

(b)(ii)

As g is analytic on \mathbb{C} and $g'(i) \neq 0$ then a small disc centred at i is mapped approximately (A4, Sect. 1, Para. 11) to a small disc centred at $g(i) = -i$. The disc is rotated by $\text{Arg}(g'(i)) = \text{Arg}(-3) = \pi$, and scaled by a factor $|g'(i)| = 3$.

(b)(iii) Γ_1 is the circle and Γ_2 the vertical line.(b)(iv) The horizontal line in the diagram below is $g(\Gamma_1)$. (A4, Sect. 4, Para. 4)

(b)(v)

Let Γ_3 be the smooth path given by $\Gamma_3 : \gamma_3(t) = t$ ($t \in \mathbb{R}$).

Γ_2 and Γ_3 meet at a right-angle at $z = 0$.

For $i = 2, 3$ we have $(g \circ \gamma_i)'(t) = \gamma_i'(t)g'(\gamma_i(t)) = \gamma_i'(t)3(\gamma_i(t))^2$. So $(g \circ \gamma_i)'(0) = 0$.

As the images of Γ_2 and Γ_3 do not meet at right-angles at 0 then g is not conformal at 0.

Question 10

(a)

$$\sin z = z \left(1 - \frac{z^2}{3!} + \frac{z^4}{5!} - \dots \right), \text{ for } z \in \mathbb{C}. \quad (\text{B3, Sect. 3, Para. 5})$$

$$\begin{aligned} \frac{z}{\sin z} &= \left(1 - \frac{z^2}{3!} + \frac{z^4}{5!} - \dots \right)^{-1} \\ &= 1 + \left(\frac{z^2}{3!} - \frac{z^4}{5!} + \dots \right) + \left(\frac{z^2}{3!} - \frac{z^4}{5!} + \dots \right)^2 + \dots \\ &= 1 + \frac{z^2}{6} + z^4 \left(-\frac{1}{120} + \frac{1}{36} \right) + \dots \end{aligned}$$

Therefore the Laurent series about 0 for f is $1 + \frac{z^2}{6} + \frac{7}{360}z^4 + \dots$ for $0 < |z| < \pi$

$\frac{1}{z^2 \sin z} = \frac{f(z)}{z^3}$ is analytic on the punctured disc $\mathbb{C} - \{0\}$. The Laurent series about 0 is

$$\frac{1}{z^3} + \frac{1}{6z} + \frac{7}{360}z + \dots = \sum_{n=-\infty}^{\infty} a_n z^n$$

As C is a circle with centre 0 then (B4, Sect. 4, Para. 2)

$$\int_C \frac{1}{z^2 \sin z} dz = 2\pi i a_{-1} = 2\pi i(2) = 4\pi i.$$

(b)

The domain of $A = \mathbb{C} - \{n\pi : n \in \mathbb{Z}\}$.

Suppose that g is another analytic function with domain A which agrees with f on $\{iy : y > 0\}$

The set $S = \left\{ i\left(1 + \frac{1}{n}\right) : n = 1, 2, 3, \dots \right\} \subseteq A$ and has the limit point $i \in A$.

f agrees with g throughout the set $S \subseteq A$ and S has a limit point which is in A . By the Uniqueness theorem (B3, Sect. 5, Para. 7) f agrees with g throughout A . Hence f is the only analytic function with domain A such that $f(iy) = \frac{y}{\sinh y}$ for $y > 0$.

(c) 6 marks

Since $\sin z = 0$ when $z = 0, z = \pm\pi, z = \pm 2\pi, \dots$. Then $f(z)$ has singularities of the form $k\pi, k \in \mathbb{Z}$.

Singularity at $z = 0$.

At $z = 0$ we can use the Laurent series found in part (a). Since $f(0) = 1$ then the singularity at 0 is a removable singularity.

Singularities at $z = k\pi$ where $k \in \mathbb{Z} - \{0\}$.

$$\sin(z - k\pi) = \sin z \cdot \cos k\pi - \cos z \cdot \sin k\pi = (-1)^k \sin z.$$

$$\text{Therefore } f(z) = \frac{z}{\sin z} = (-1)^k \frac{z}{\sin(z - k\pi)} = (-1)^k \left\{ \frac{z - k\pi}{\sin(z - k\pi)} + \frac{k\pi}{\sin(z - k\pi)} \right\}$$

$$\lim_{z \rightarrow k\pi} \frac{z - k\pi}{\sin(z - k\pi)} = 1$$

As $\lim_{z \rightarrow k\pi} (z - k\pi) \frac{k\pi}{\sin(z - k\pi)} = k\pi$ then there is a simple pole at $z = k\pi$ (B4, Sect. 3, Para. 2).

Therefore there f has simple poles at $k\pi$ where $k \in \mathbb{Z} - \{0\}$.

Question 11

(a)

Let $f(z) = \exp(z^4)$ and $R = \{z : |z| < 2\}$.

As f is analytic on the bounded region R and continuous on \bar{R} then by the Maximum Principle (C2, Sect. 4, Para. 4) there exists an $\alpha \in \partial R = \{z : |z| = 2\}$ such that $|f(z)| \leq |f(\alpha)|$ for $z \in \bar{R}$.

When $z \in \partial R$ we can write it in the form $z = 2 \exp(i\theta)$, where θ is real.

Then $|\exp(z^4)| = \exp(\operatorname{Re}(z^4)) = \exp(16 \cos 4\theta)$.

This is a maximum when 4θ is a multiple of 2π and it equals e^{16} .

Therefore $\max \{ |\exp(z^4)| : |z| \leq 2 \} = e^{16}$. The maximum is attained when $z = 2e^0 = 2$, $z = 2e^{\pi/2} = 2i$, $z = 2e^\pi = -2$, and $z = 2e^{3\pi/4} = -2i$.

(b)

Let $D_f = \{z : |z| < 5\}$ and $D_g = \{z : |z| > 5\}$.

Since $D_f \cap D_g = \emptyset$ then f and g are not direct analytic continuations of each other.

Let $h(z) = \frac{1}{1 - \frac{z}{5}} = \frac{5}{5 - z}$ on D_h , where $D_h = \mathbf{C} - \{5\}$.

When $z \in D_f$ then $|z|/5 < 1$ and the geometric series $\sum_{n=0}^{\infty} \left(\frac{z}{5}\right)^n$ is convergent and has the sum

$$\frac{1}{1 - \frac{z}{5}} = \frac{5}{5 - z}. \quad (\text{B3, Sect. 3, Para. 5})$$

Since $f = h$ when $z \in D_f \subseteq D_f \cap D_h$ then h is an analytic continuation of f .

When $z \in D_g$ then $5/|z| < 1$ and the geometric series $\sum_{n=0}^{\infty} \left(\frac{5}{z}\right)^n$ is convergent and has the sum

$$\frac{1}{1 - \frac{5}{z}} = \frac{z}{z - 5}.$$

Therefore $-\sum_{n=1}^{\infty} \left(\frac{5}{z}\right)^n = -\frac{5}{z} \sum_{n=0}^{\infty} \left(\frac{5}{z}\right)^n = -\frac{5}{z-5}$ when $z \in D_g$.

Since $g = h$ when $z \in D_g \subseteq D_g \cap D_h$ then g is an analytic continuation of h .

Since (f, D_f) , (g, D_g) , (h, D_h) form a chain then f and g are indirect analytic continuations of each other.


Question 12

(a)

Using the formula for a transformation mapping points to the standard triple (D1, Sect. 2, Para. 11) then the Möbius transformation \hat{f}_1 which maps i , ∞ , and $-i$ to the standard triple of points 0 , 1 , and ∞ respectively is

$$f_1(z) = \frac{(z-i)(\infty+i)}{(z+i)(\infty-i)} = \frac{z-i}{z+i}$$

(b)(i)



(b)(ii) Since \hat{f}_1 maps i to 0 and $-i$ to ∞ then the curved boundaries of R are mapped to extended lines originating at the origin. As the angle between the arcs of the circles meeting at i is $\pi/2$ and the mapping \hat{f}_1 is conformal then the angle between the extended lines at the origin is also $\pi/2$.

Points in R , which lie on the imaginary axis, can be written as

$$z = i(1 - \delta) \quad \text{where } 0 < \delta < 2.$$

For these points we have $f_1(z) = \frac{i(1-\delta-1)}{i(2-\delta)} = -\frac{\delta}{2-\delta}$.

Therefore \hat{f}_1 maps these points to the negative real-axis. In R this line is at an angle of $\pi/4$ to the two arcs of the circles so, as it is a conformal mapping, then in the image of R this is also true.

Therefore the image of R under \hat{f}_1 is R_1 is S .

(b)(iii) A conformal mapping from S onto T is the power function $w = z^2$.

Since the combination of conformal mapping is also conformal then a conformal mapping from R to T is

$$f(z) = \left(\frac{z-i}{z+i} \right)^2.$$

(b)(iv)

D1, Sect. 4, Para. 5 gives

$$h(z) = \frac{z-1}{z+1}$$

as a conformal mapping from T onto the open unit disc.

Therefore a one-one conformal mapping from R onto the open unit disc is

$$g(z) = \frac{\left(\frac{z-i}{z+i} \right)^2 - 1}{\left(\frac{z-i}{z+i} \right)^2 + 1}.$$